# A review of the epidemiology of epilepsy in Mexico during 1970 to 2020

# Spanish title: Revision sistematica de la epidemiologia en Mexico de la epilepsia de

1970 a 2020.

# A systematic review

### Short title: Epidemiology of epilepsy in Mexico

Gerardo Quiñones Pesqueira, M.D. ID: https://orcid.org/0000-0003-4420-5107 a						
Daniel San Juan, M.D M.Sc. <sup>b</sup> ht	tps://orcid.org/ 0000-0001-6685-5851					
Rosana Huerta Albarra, M.D. <sup>c.</sup>	https://orcid.org/ 0000-0003-4231-3358					
Maximo Leon Vazquez, M.D. <sup>d</sup>	https://orcid.org/0000-0002-7213-7837					
Gerardo Quiñones Canales, M.D. e	https://orcid.org/0000-0003-4445-5683					
Jorge Gonzalez Pesqueira <sup>f.</sup>	https://orcid.org/0000-0002-4297-1329					

<sup>a</sup>Department of Neurophysiology, National Institute of Neurology and Neurosurgery Manuel Velasco Suárez,

Mexico City, Mexico.

<sup>b</sup>Epilepsy Service, National Institute of Neurology and Neurosurgery Manuel Velasco Suárez, Mexico City,

Mexico.

<sup>c</sup>Department of Neuropediatrics, Hospital General de México Dr Eduardo Liceaga, Mexico City

<sup>d</sup> Department of Neuroscience, Insituto Mexicano del Seguro Social Centro Médico Nacional La Raza, Mexico

City, Mexico

<sup>e</sup> Department of Neuroscience, Instituto de Seguridad y Servicios Sociales de los Trabajadores del Estado,

Durango, Mexico.

<sup>f</sup> University of Arizona, College of Medicine-Phoenix, United State of America.

Correspondence: Daniel San-Juan, MSc. Epilepsy Service, National Institute of Neurology and Neurosurgery.

Insurgentes Sur 3877, La Fama, Tlalpan, Mexico City, Mexico Zip code 14269. Tel.: +52 5556063822 E-mail address:

dsanjuan@innn.edu.mx

Declarations of interest: None.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-

profit sectors.

Authors' contributions: All authors worked extensively on the manuscript, and all approved this submitted version. The authors Gerardo Quiñones Pesqueira and Daniel San Juan conceived the idea for the article, which was written by Gerardo Quiñones Pesqueira and critically reviewed by all authors.

### Abstract:

**Background:** Epilepsy is the most common major neurological disorder that affects people of all ages. The prevalence varies from one country to another and even between different areas, due to a lack of access to medical care for reasons related to limited resources.

**Objective:** Epilepsy is a worldwide public health problem that affects more deeply populations living in developing countries such as Mexico, where more aggressive health policies based on epidemiological data are needed, however, this information is scarce and the evolution of this data over time remains unclear. The aim of this study is to provide an overview of the epidemiology of epilepsy in Mexico (1970-2020).

**Methods:** We searched descriptive epidemiological studies on epilepsy in rural and urban regions of Mexico from 1970 to 2020. Available data on the socio-demographic characteristics, prevalence, and incidence data were extracted. Finally, the abstract, full-text review, and data abstraction were conducted in duplicate and reported using PRISMA. Descriptive statistics was also used.

**Results:** Overall, 11 underrepresented and heterogeneous epidemiological studies were included. In total, the prevalence of epilepsy in Mexico is 3.9-41:1000 inhabitants; 3.9 to 41 per 1,000 persons in rural regions and 3.49 to 44.3 per 1,000 persons in urban regions. None of these studies addressed the incidence of epilepsy. The prevalence of epilepsy in Mexico has remained unchanged during the last five decades.

**Significance:** Our results confirm a high prevalence of epilepsy in both urban and rural settings in Mexico that remain unchanged during the last five decades. All studies included in this review showed multiple methodological limitations. New and robust epidemiological studies are needed to delineate the epidemiological profile of epilepsy in Mexico.

Keywords: Epilepsy; Epidemiology; Prevalence Mexico; Rural; Urban; Developing country

#### **Resumen:**

La epilepsia es el trastorno neurológico más común que afecta a individuos de todas las edades. La prevalencia varía entre paises e incluso entre diferentes áreas del mismo pais debido a la falta de acceso a la atención médica. **Objectivo:** La epilpesía es un problema de salud pública mundial que afecta de manera más importante a los países en desarrollo como México, donde hacen falta políticas en salud y datos epidemiológicos, para conocer el impacto real de esta enfermedad. El objetivo de este estudio es brindar una visión general de la epidemiología de la epilepsia en México (1970-2020). Métodos: Se realizo unas búsquedas bibliografica de estudios epidemiológicos descriptivos sobre epilepsia en regiones rurales y urbanas de México desde 1970 a 2020. Se extrajeron los datos disponibles sobre las características sociodemográficas, prevalencia e incidencia. Finalmente, el resumen, la revisión completa del texto y la extracción de datos se realizaron por duplicado y se informaron utilizando PRISMA. Se utilizo estadística descriptiva. Resultados: Se incluyeron 11 estudios epidemiológicos heterogeéneos. En total la prevalencia de epilepsia en México es 3.9-41: 1000 habitantes; 3,9 a 41 por 1000 personas en las regiones rurales y 3,49 a 44,3 por 1000 personas en las regiones urbanas. Ninguno de estos estudios abordó la incidencia de epilepsia. La prevalencia de la epilepsia en México se ha mantenido sin cambios durante las últimas cinco décadas. Importancia: Nuestros resultados confirman una alta prevalencia de epilepsia en las areas urbanas y rurales de México que se mantiene sin cambios durante las últimas cinco décadas. Todos los estudios incluidos en esta revisión mostraron múltiples limitaciones metodológicas. Se necesitan estudios epidemiológicos nuevos y sólidos para delinear el perfil epidemiológico de la epilepsia en México. Palabras claves: Epilepsia Epidemiología; Prevalencia en Mexico; Rural; Urbana; Paises en desarrollo.

# Introduction

Epilepsy is the most common major neurological disorder that affects people of all ages. Worldwide the frequency of epilepsy is estimated to be 69 million people, with a prevalence of active epilepsy at 6.38 per 1000 inhabitants<sup>1</sup>. Forty-five (65%) out of 69 million people with epilepsy live within rural regions in developing countries, 17 (25%) million are living in the urban areas of these countries, and the rest, 7 (10%) million people, are living in developed countries<sup>2</sup>. About 5 million live in Latin America and the Caribbean, with a median prevalence interval ranging from 15.8 to 17.8 patients per 1000 inhabitants<sup>3</sup>. Incidence rates are higher in developing countries especially in Africa and Latin America, where figures can exceed 150 per 100,000 inhabitants<sup>3</sup>. Globally, almost 2.4 million people are diagnosed with epilepsy each year<sup>5</sup>.

The prevalence of epilepsy varies from one country to another and even between different areas within the same country due to a lack of access to medical care for reasons related to limited resources, geography, and due to problems with communicating epidemiological data and addressing it appropriately. To this point less than 40% of Latin American countries had at least one epidemiological study on epilepsy. In fact, there is sparce epidemiological information available from local health authorities in these countries, and few epidemiological studies have been conducted, which are essential for creating plans and strategies to address the disparities of epilepsy cases at the local level as recommended by The PanAmerican Health Organization (PAHO) as noted in the last report released in 2013<sup>12</sup>.

Mexico is a developing country located in Latin America with more than 126 million inhabitants (78.8% urban and 21.2% rural)<sup>14</sup>, however, like other developing countries, there

has been a lack of overview of the epidemiological studies conducted in the last decades which are necessary for the support of public health policies and strategies in reducing the gaps in the number of epilepsy cases, especially in areas with a higher burden of disease.

*Demographic data of Mexico*. Mexico is located in North America and covers 1,972,550 km<sup>2</sup>, with a density of about 57 people per km<sup>2</sup><sup>6</sup>. Acording to The National Institute of Statistics and Geography (INEGI) survey 2020 in Mexico, the population of Mexico is 126,014,024 inhabitants, with a total of 78.84% living in urban areas and 21.16% living in rural areas; 48.8% men and 51.2% women<sup>6</sup>. The distribution in percentages of the inhabitants for the age groups is 18.6% (0- 14 years old), 67.3% (15- 64 years old) and 14.1% (65 or over) <sup>14</sup>.

The aim of our review is to provide an outline of the epidemiological studies of epilepsy in both urban and rural regions of Mexico from 1970 to 2020.

# **Materials and Methods**

We conducted this review adhered to the methods described in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) reporting guidelines<sup>15.</sup>

#### Search strategy

We searched between November and December 2020 for epidemiological studies (1970-2015) in Spanish or English conducted in Mexico in five databases: Pubmed, Scielo, LILACS (Health Information from Latin America and the Caribbean), Medigraphic, and OVID. The

mesh terms used were: «epilepsy», «Mexico», «Latin America», «epidemiology», «prevalence», «incidence», and «distribution». The exactitude of the search strategy was validated by cross-verification with the results of previous reviews. Likewise, we reviewed the bibliographic reference lists of the included studies and previous systematic and descriptive reviews on that subject.

#### Study selection

Abstracts and titles of all references were screened in duplicate by 2 independent reviewers to identify original population-based studies on the prevalence or incidence of epilepsy in Mexico.

Articles were included if they met the following inclusion criteria: (1) population-based studies that provide a standardized definition of epilepsy having more than one stage of diagnosis and (2) that the main objective of the study is to determine the prevalence and/or incidence of epilepsy in adults and/or children in Mexico without any language restrictions. 3) reported a prevalence or incidence of epilepsy. We excluded descriptive case-series from single centers, review studies, or other type of studies of epilepsy in Mexico. Disagreements pertaining to the inclusion of articles were resolved by consensus or involvement of a third author as necessary.

#### Data extraction

Data abstraction was completed in duplicated by 2 independent reviewers using a standardized data collection form and stored in an Excel sheet. Variables extracted were author, year of publication, socio-demographic characteristics, location, type of population

6

area (urban or rural), type of study, number of patients, prevalence of epilepsy, and age group (only children, only adults or both). Regarding age, we stratified the cases into 3 large groups for age-specific distribution: 1. Children (preschool and school age); 2. Adults 18 years and older; 3. Any age (children and adults). For the definition of epilepsy, we relied on a common definition which defines epilepsy as two or more unprovoked seizures more than 24 h apart. We defined prevalence as the number of existing cases of epilepsy in a population over the total population at a specific point in time. In this study, we focused on two types of prevalence: the lifetime prevalence of epilepsy (including active and inactive epilepsy/historical cases over time between birth and assessment) and (2) the prevalence of active epilepsy (individuals reporting recent seizures or those currently taking anticonvulsants). The populational study was defined as a study of a group of individuals taken from the general population who shares a common characteristic, such as age, sex, or health condition<sup>16</sup>.R egarding incidence, none of these studies addressed incidence of epilepsy. After reviewing the abstracts of the articles found; 11 studies were chosen focused on the prevalence of epilepsy in Mexico.

#### Data synthesis and analysis

We used descriptive statistics to calculate frequencies, percentages, and the rates of both prevalence and incidence of epilepsy in Mexico based on 1000 inhabitants. This was done using Excel, Microsoft Software version 2020.

## Results

Initially, there were 220 scientific articles recognized, 170 were deemed out-ofscope of the epidemiological study design and were excluded solely based on their abstract. Then, out of the remaining papers, 15 were selected for full text review, and subsequently, 11 scientific articles met the study criteria and were selected for this review. Figure 1. Tables 1 and 2 show the epidemiological studies according to the rural or urban areas covered. Six (54.4%) studies were carried out in urban areas and five (45.6%) studies in rural communities. There was no disagreement among reviewers. The epidemiological studies analyzed where methodological heterogenous and included few urban and rural regions of Mexico as is shown in Figure 2.

In total, the prevalence of epilepsy (urban and rural) was 3.9-41:1000 inhabitants. We found a prevalence of epilepsy ranging from 3.9 to 41 per 1,000 persons in rural regions and 3.49 to 44.3 per 1,000 persons in urban regions of Mexico. None of these studies addressed incidence of epilepsy. The prevalence of epilepsy in the epidemiological studies analyzed remain unchanged in rural and urban settings during the last five decades in Mexico.

## Discussion

This systematic review found a high prevalence of epilepsy in urban and rural regions in Mexico during the last five decades compared to developed countries and globally <sup>17</sup>. The prevalence of epilepsy in the epidemiological studies analyzed remain unchanged in rural and urban settings during the last five decades in Mexico. However, there was a significant heterogeneity between the epidemiological studies and caution is needed in interpreting these results. The current etiology of epilepsy in Latin America and Caribbean (LAC continues to be associated with neurocysticercosis (NCC) in 17.3% of the epidemiological studies in this region. Unfortunately, the etiology in Mexican epidemiological studies has not been addressed appropriately, however, local studies show that NCC is considered one of the leading causes of epilepsy<sup>31</sup>.

Mexico is a middle-income country, with 62.25 million people living in poverty, which may explain the higher risk of developing epilepsy given that it is reported to a greater degree in the sector of the population considered to occupy a lower socioeconomic status. Furthermore, no epidemiological studies concerning incidence of epilepsy in Mexico were found. Finally, low levels of education have been shown to translate into low-income levels which are related to poor health conditions, a likely factor for the higher incidence of epilepsy in Mexico<sup>33</sup>.

The WHO's 2016 Global Burden of Disease study of epilepsy reported an estimation of 45.9 million patients with all-active epilepsy globally, accounting 0.56% of total disability-adjusted life-years (DALYs) internationally. Also, found a greater severity and higher sum of years of life lost for premature mortality and years lived with disability due epilepsy in low-income settings<sup>17</sup>. The prevalence of epilepsy worldwide differs significantly among countries depending on the sociodemographic characteristics, as well as both, the local distribution of risk and etiologic factors. The overall global prevalence of epilepsy was 7.60 per 1,000 inhabitants (95% CI 6.17–9.38)<sup>35</sup>. According to a recent systematic review and meta-analysis of population-based studies of the epidemiology of epilepsy in LAC, there is a lifetime prevalence that ranges from 11.7 to 16.6 patients per 1000 inhabitants with an incidence range of 111.24 per 100 000 person-years (95% CI = 64.88–169.51)<sup>36</sup>. Our results

are included in this data. Currently, Mexico is considered one of the countries with the highest prevalence of epilepsy (25:1000) in Latin America followed by Chile (17.75:1000) and Guatemala (12.95:1000), however, this estimation was established based on only four epidemiological studies from Mexico<sup>36</sup>. Interestingly, in this study there were no differences found in relation to setting, whether it be urbal/rural, or differences attributable to sex or age clusters, or even the income in LAC.

A recent report of the Prevention Task Force of the International League of Epilepsy conducted a systematic review of published epidemiologic studies of epilepsy of 4 preventable etiologic categories: perinatal insults, traumatic brain injury (TBI), CNS infection and stroke. In fact, CNS infections were a more common attributed cause in Lowerand Middle-income countries (LMIC), accounting for about 5% of all epilepsy cases. Among some rural LMIC communities, the median proportion of epilepsy cases attributable to endemic NCC was as high as 34%. Therefore, a large proportion of the overall public health burden of epilepsy is due to preventable causes as well as the attributable fraction for perinatal causes, infections, TBI and stroke, in sum reaching nearly 25%<sup>43</sup>.

Significant limitations of the epidemiological studies used for this review include the low sample size of each study, the heterogeneity of the studies, insufficient detail in describing the epilepsy types, the absence of stating epilepsy status (active or resolved), and the lack of mention of etiological sources.

Worthy of attention are the results of our study that show that the prevalence of epilepsy in Mexico in both rural and urban settings remain unchanged during the last five decades, even with scientific and technological advances in the sector of corresponding health issues, all the while, there continues to be a lack of access to health services in developing countries, and even more, in rural communities which greatly limit the diagnosis and treatment of epilepsy. This causes the prevalence to be higher in these regions compared to developed countries or communities with greater access to health services<sup>44</sup>. In such a way that epilepsy is a public health problem that must be taken seriously, public policies must reflect this and be generated in order to guarantee access to adequate health services.

Furthermore, new, robust, and broad prospective epidemiological studies are needed to determine the national epidemiological profile of patients with epilepsy in Mexico to contribute delineated health policies, research investment and dedicated health infrastructure to reduce the existing gap disparities in epilepsy.

#### **Conclusions:**

Our results confirm a high prevalence of epilepsy in both urban and rural settings in Mexico that remain unchanged during the last five decades. All studies included in this review showed multiple methodological limitations. The heterogeneity of the results could be due to differences in the population samples, the use of differing screening methods, the definitions and classifications used in the questionnaires, and selection bias. Caution is needed to interpret our findings. New and robust epidemiological studies are needed to delineate the epidemiological profile of epilepsy in Mexico. 'We confirm that we have read the Journal's position on issues involved in

ethical publication and affirm that this report is consistent with those

guidelines.'

# References

Journal article:

1. World Health Organization. Epilepsy. Fact sheet. No. 999. 2012. Updated February 2016. Available in: http://www.who. int/mediacentre/factsheets/fs999/en/

2. Ngugi AK, Bottomley C, Kleinschmidt I, et al. Estimation of the burden of active and life-time epilepsy: a meta-analytic approach. Epilepsia. 2010;51(5):883–890.

3. Meyer AC, Dua T, Ma J, et al. Global disparities in the epilepsy treatment gap: a systematic review. Bull World Health Organ. 2010;88(4):260–266. doi: 10.2471/BLT.09.064147.Epub 2009 Sep 25.

4. Gracia F. Epidemiología de las epilepsias en La noamérica. En: Medina MT, Chaves-Sell F, Chinchilla N, Gracia F, editores. Epilepsias en Centroamérica. Tegucigalpa: Scancolor;

2001 [consultado el 18 de agosto del 2013]; p. 17-22.

5. Dévora Keste, Carlos Acevedo, Medina MT, Tomás Mesa, Jorge Rodríguez; Epilepsy in Latin America; Technical document based on presentations at the international work shop in Santiago, Chile, in August 2013. https://www.ilae.org/files/dmfile/PAHO-report2016-English1.pdf

6. "Numeralia indígena 2015", en Indicadores Socioeconómicos de los Pueblos Indígenas de México, 2015", CDI, México, 2015,

disponibleen: https://www.mx/inpi/articulos/indicadores-socioeconomicos-de-los-pueblos-indige- nas-de-mexico-2015-116128.

7. Burneo JG, Tellez-Zenteno J, Wiebe S. Understanding the burden of epilepsy in Latin America: a systematic review of its prevalence and incidence. Epilepsy Res. 2005; 66: 63-74. DOI: https://doi.org/10.1016/j.eplepsyres.2005.07.002

8. OMS (2015). Carga mundial de epilepsia y necesidad de medidas coordinadas en los países para abordar sus consecuencias sanitarias y sociales y su conocimiento por el público (A68/12) (Resolución WHA68.20). Ginebra: OMS; 2015. Disponible en:

http://apps.who.int/gb/ebwha/pdf\_files/WHA68/A68\_12-sp.pdf

http://apps.who.int/gb/ebwha/pdf\_files/WHA68/A68\_R20-sp.pdf

9. Caraveo-Andagua, J., Medina-Mora, M.E., et al., 1996. Prevalence of psychiatric

dysfunction in the urban population of Mexico. Salud. Ment. 19, 14-21.

10. García-Pedroza F, Millán R, Peñaloza Y. Epidemiología clinica de la epilepsia. Rev Mex Neuroci 2010;11:82–102.

11. Beghi E, Hesdorffer D. Prevalence of epilepsy--an unknown quantity. Epilepsia. 2014 Jul;55(7):963-7. doi: 10.1111/epi.12579. Epub 2014 Jun 25. PMID: 24966095.

12. Organización Panamericana de la Salud, Organización Mundial de la Salud. Liga Internacional Contra la Epilepsia–Buró Internacional para la Epilepsia. Estrategia y Plan de Acción sobre la Epilepsia. Washington (DC), Estados Unidos:

OPS;2011[http://www.paho.org/hq/dmdocuments/epilepsia%20doc.pdf,last accessed on March 2015].

13. Barendregt JJ, Doi SA, Lee YY, Norman RE, Vos T. Meta- analysis of prevalence. J Epidemiol Commun Health. 2013;67:974.

14. All-Mexican Population Census, 2020Instituto Nacional de Estadistica y Geografia;Inegi.org.mx .

15. Hamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P):elaboration and explanation. BMJ. 2016 Jul 21;354:i4086.https://doi.org/10.1136/bmj.i4086

16. Banerjee, Amitav & Chaudhury, Suprakash. (2010). Statistics without tears: Populations and samples. Industrial psychiatry journal. 19. 60-5. 10.4103/0972-6748.77642.

17. GBD 2016 Epilepsy Collaborators. Global, regional, and national burden of epilepsy, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Neurol. 2019 Apr;18(4):357-375. doi: 10.1016/S1474-4422(18)30454-X. Epub 2019 Feb 14. Erratum in: Lancet Neurol. 2019 May;18(5):e4. PMID: 30773428; PMCID: PMC6416168.

18. Senanayake N, Román GC. Epidemiology of epilepsy in developing countries. Bull World Health Organ. 1993;71(2):247-58. PMID: 8490989; PMCID: PMC2393447.

19. Márquez L, Olivares L. Epilepsia en México: estudio epidemiológico de una comunidad rural. Salud Publica Mex. 1979; 21 (5): 487-495.

20. Gutierrez Avila, J.H., 1980. Prevalence of epileptic seizures among children in a rural town of Mexico. Salud. Publica. Mex. 22, 269–273.

21. Garcia-Pedroza, F., Rubio-Donnadieu, F., et al., 1983. Prevalence of epilepsy in children: Tlalpan Mexico City, Mexico. Neuroepi- demiology 2, 16–23.

22. Quet F, Preux PM, Huerta M, Ramirez R, Abad T, Fragoso G, et al. Determining the burden of neurological disorders in populations living in tropical areas: who would be questioned? Lessons from a Mexican rural community. Neuroepidemiology. 2011;36(3):194-203. doi: 10.1159/000327496.

23. San-Juan D, Alvarado-León S, Barraza-Díaz J, Davila-Avila NM, Hernandez-Ruíz A, Anschel DJ. Prevalence of epilepsy, beliefs and attitudes in a rural community in Mexico: a door- to-door survey. Epilepsy Behav. 2015; 46: 140-143.

24. Cruz-Alcala, L.E., Velazquez-Castellanos, J.L., 2002. Prevalencia de algunas enfermedades neurologicas en la ciudad de Tepatitlan Jalisco Mexico. Rev. Mex. Neuroci. 3, 71–76.

25. Gutierrez-Avila, H., Rubio-Donnadieu, F., et al., 1980. Prevalence of epilepsy in school-age children of an urban community in Mexico City. Gac. Med. Mex. 116, 497–501
26. Olivares L. La epidemiología neurológica en México. Salud Publica Mex 1976;18:665-672.

27. García-Pedroza F, Rubio-Donnadieu F. Prevalencia de las crisis epilépticas en 23.000 niños de 3 año de primaria en escuelas públicas de 30 estados de la República Mexicana [Prevalence of epileptic crises in 23,000 third-year primary school children in public schools in 30 Mexican states]. Epilepsia 1991;32 (suppl 1):111.

28. García Pedroza F, Rubio-Donnadieu F. Prevalencia de la Epilepsia en Naolinco, Veracruz [Prevalence of Epilepsy in Naolinco, Veracruz]. Archivos INNN 1993;vol.8(3):168.

29. Beghi, E. (2019). The Epidemiology of Epilepsy. Neuroepidemiology, 1 7.doi:10.1159/000503831.

30. Bruno E, Bartoloni A, Zammarchi L, Strohmeyer M, et al. Epilepsy and neurocysticercosis in Latin America: a systematic review and meta-analysis. PLoS Negl Trop Dis. 2013;7:e2480.

31. Fleury A, Sciutto E, Larralde C. Neurocysticercosis is still prevalent in Mexico. Salud Publica Mex. 2012 Nov-Dec;54(6):632-6. doi: 10.1590/s0036-36342012000600012. PMID: 23318900.

32. Banerjee PN, Filippi D, Allen Hauser W. The descriptive epidemiology of epilepsy-a review. Epilepsy Res 2009;85: 31–45.

33. Programa Especial de los Pueblos Indígenas 2014-2018, Primera edición, 2014 http://www.cdi.gob.mx

34. Murray CJ, Vos T, Lozano R, et al. . Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2197–2223

35. Fiest KM, Sauro KM, Wiebe S, Patten SB, Kwon CS, Dykeman J et al. Prevalence and incidence of epilepsy: A systematic review and meta-analysis of international studies. Neurology. 2017 Jan 17;88(3):296-303. doi: 10.1212/WNL.000000000003509. Epub 2016 Dec 16. Erratum in: Neurology. 2017 Aug 8;89(6):642. PMID: 27986877; PMCID: PMC5272794

36. Alva-Díaz C, Navarro-Flores A, Rivera-Torrejon O, Huerta-Rosario A, et al. Prevalence and incidence of epilepsy in Latin America and the Caribbean: A systematic review and meta-analysis of population-based studies. Epilepsia. 2021 Apr;62(4):984-996. doi: 10.1111/epi.16850.

37. Giussani G, Canelli V, Bianchi E, Erba G, Franchi C, Nobili A, et al.; EPIRES Group. Long-term prognosis of epilepsy, prognostic patterns and drug resistance: a population-based study. Eur J Neurol. 2016 Jul;23(7):1218–27

38.Mbuba CK, Ngugi AK, Newton CR, Carter JA. The epilepsy treatment gap in developing countries: a systematic review of the magnitude, causes, and intervention strategies. Epilepsia. 2008 Sep;49(9):1491–503.

39.Watts AE. The natural history of untreated epilepsy in a rural community in Africa. Epilepsia. 1992 May-Jun;33(3):464–8.

40.Senanayake N, Román GC. Epidemiology of epilepsy in developing countries. Bull World Health Organ. 1993;71(2):247–58.

41"Poverty in Mexico 2018. Población según pertenencia étnica", CONEVAL, México,2018,en: <u>https://www.coneval.org.mx/Evaluacion/PublishinlgImages/Pobreza\_201</u> 8/POBLACION PERTENENCIA ETINICA.jpg.

42 "Plan Nacional de Desarrollo 2019-2024". SEGOB, México, 12 de julio de 2019. https://www.gob.mx/cenace/acciones-y-programas/plan-nacional-de-desarrollo-2019-2024-195029.

43. Thurman DJ, Begley CE, Carpio A, Helmers S, Hesdorffer DC, Mu J, et al. The primary prevention of epilepsy: A report of the Prevention Task Force of the International League Against Epilepsy. Epilepsia. 2018 May;59(5):905-914. doi: 10.1111/epi.14068. Epub 2018 Apr 10. PMID: 29637551; PMCID: PMC7004820

44. Thurman DJ, Logroscino G, Beghi E, Hauser WA, et al.; Epidemiology Commission of the International League Against Epilepsy. The burden of premature mortality of epilepsy in high-income countries: A systematic review from the Mortality Task Force of the International League Against Epilepsy. Epilepsia. 2017 Jan;58(1):17-26. doi: 10.1111/epi.13604. Epub 2016 Nov 26. PMID: 27888514; PMCID: PMC7004822.

Figure 1. Flowchart of epidemiological studies related to epilepsy in Mexico.





**Figure 2.** Geographical distribution of Mexico where epilepsy prevalence studies were carried out.

Author	Study	Location	Population	n	Prevalence	Age group
	design	and setting	area		x 1000	
Olivares-L	Case-	Mexico City	Urban	2,169	3.49	Children and
1976 <sup>26</sup> .	Control	Third level				adults
		center				
H Gutíerrez-	Cross- Sectional	Mexico City	Urban	1,013	16	School-age
Avila	Study	Primary			12.6- 23.6	ernieren
1980 <sup>25</sup>		schools				
Garcia-	Cross-	Tlalpan	Urban	1,934	44.3	
Pedroza	Study	Mexico City				Children
1983 <sup>21</sup>						
Garcia-	Cross- Sectional	Mexico City	Urban	23,000	10.8	Children
Pedroza	Study					
1991 <sup>27</sup>						
Garcia-	Door-to-	Comalcalco,	Urban	142,000	20	Children and
Pedroza	door survey	Tabasco				adults
1993-1994						
Cruz-Alcala	Door-to-	Tepatlitan,	Urban	9,082	6.8	Children and
2002 <sup>24</sup>	door survey	Jalisco				adults

**Table 1.** Epidemiological studies on the prevalence of epilepsy in urban areas ofMexico (1970-2015).

Study	Design	Location	Population	n	Prevalence	Evaluated
			Area		x 1000	group
Marquez-Olivares	Door-	Luvianos	Rural	4103	5.8	Children
1979 <sup>19</sup>	to-door	Tejupilco				and adults
	survey	State of				
		Mexico				
Gutiérrez-A.	Door-	San Miguel	Rural	360	25-41	Children
1980 <sup>20</sup>	to-door	Tecomatlàn				6-12 years-
	survey	State of				old
		Mexico				
Gutiérrez-A.	Door-	Nanolinco,	Rural	2,025	11	School-age
1992 <sup>28</sup>	to-door	Veracruz				Children
	survey					
Quet	Door-	San Andres	Rural	4,008/6,203	3.9	Children
2011 <sup>22</sup>	to-door	Azumiatla,				and adults
	survey	Puebla				
San-Juan	Door-	Xocotitlan,	Rural	863	25.4	Children
2015 <sup>23</sup>	to-door	Hidalgo				and adults
	survey					

**Table 2.** Studies on the prevalence of epilepsy in rural areas of Mexico (1970-2015).